

# LESSON 12

## THREE DIMENSIONAL GEOMETRY

### 1. INTRODUCTION


To locate a point in a plane we need a two dimensional coordinate system. On similar lines to locate a point in space we use a three dimensional coordinate system.

Consider three planes intersecting at a point  $O$ . The coordinate axes  $x$ -axis,  $y$ -axis and  $z$ -axis are shown by dark lines.


i.e., if we have an insect on the floor of a room, we can specify its position by a two dimensional coordinate system but if a spider is hanging from the roof of a room we specify its position in a three dimensional coordinate system.

### 2. COORDINATE AXES


Three dimensional coordinate system is formed by taking three mutually perpendicular planes intersecting at a point  $O$ .



The lines  $XOX'$ ,  $YOY'$  and  $ZOZ'$  at which the three planes intersect are mutually perpendicular to each other and are referred to as the  $x$ -axis,  $y$ -axis and  $z$ -axis respectively. The distance measured from  $XY$  plane upwards in the direction of  $OZ$  are taken as positive and those measured downwards in the direction of  $OZ'$  are taken as negative. We specify the coordinates of a point by specifying the  $x$ ,  $y$  and  $z$  coordinates of a point.



If the coordinates of  $P$  are  $(a, b, c)$ , we mark the coordinates of other points as shown in the figure.



### 3. DISTANCE BETWEEN TWO POINTS

To find the distance between two points  $P(x_1, y_1, z_1)$  and  $Q(x_2, y_2, z_2)$  in space.

From the above figure we note that,

$$\angle PAN = 90^\circ$$

Applying

Pythagoras theorem,

$$PN^2 = PA^2 + AN^2$$

$$\angle PNQ = 90^\circ$$

$$\therefore PQ^2 = PN^2 + NQ^2$$


$$\therefore PQ^2 = PA^2 + AN^2 + NQ^2$$

$$PA = (x_2 - x_1)$$

$$AN = (y_2 - y_1)$$

$$NQ = (z_2 - z_1)$$

$$\therefore PQ^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2$$



$$\therefore PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

**Illustration 1**

**Question:** Find the distance between the of points  $A(2, >1, 3)$  and  $B(>2, >1, 3)$ .

**Solution:** Distance =  $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

$$AB = \sqrt{(2+2)^2 + (-1+1)^2 + (3-3)^2} = 4 \text{ units}$$

**Illustration 2**

**Question:** Prove by using distance formula that the points  $P(1, 2, 3)$ ,  $Q(>1, >1, >1)$  and  $R(3, 5, 7)$  are collinear.

**Solution:** We have  $PQ = \sqrt{(-1-1)^2 + (-1-2)^2 + (-1-3)^2}$   
 $= \sqrt{4+9+16} = \sqrt{29}$

$$QR = \sqrt{(3+1)^2 + (5+1)^2 + (7+1)^2}$$

$$= \sqrt{16+36+64} = \sqrt{116} = 2\sqrt{29}$$

and  $PR = \sqrt{(3-1)^2 + (5-2)^2 + (7-3)^2} = \sqrt{4+9+16} = \sqrt{29}$

Since  $QR = PQ + PR$ . Therefore the given points are collinear.

**4. SECTION FORMULA**


To find the coordinates of a point  $R$  which divides  $PQ$  internally in the ratio  $m : n$ , drop perpendicular from  $P, R$  and  $Q$  on the  $xy$  plane. Let the foot of perpendiculars on the  $xy$  plane be denoted as  $L, N$  and  $M$ .

$\therefore \triangle PRS$  is similar to  $\triangle RQT$ .

$\therefore$  the corresponding sides are proportional.

$$\therefore \frac{PR}{RQ} = \frac{m}{n} = \frac{SP}{QT} = \frac{SR}{RT}$$

Now,  $SP = (SL - PL) = (RN - PL) = (z - z_1)$



$$QT = (QM - TM) = (QM - RN) = (z_2 - z)$$

$$\therefore \frac{m}{n} = \frac{(z - z_1)}{(z_2 - z)}$$

$$\therefore (mz_2 - mz) = nz - nz_1 \Rightarrow z(m + n) = mz_2 + nz_1$$

$$z = \frac{(mz_2 + nz_1)}{(m + n)}$$

Similarly,

$$y = \frac{(my_2 + ny_1)}{(m + n)}, \quad x = \frac{(mx_2 + nx_1)}{(m + n)}$$

\(\therefore\) The coordinates of point  $R$  are

$$\left( \frac{mx_2 + nx_1}{m + n}, \frac{my_2 + ny_1}{m + n}, \frac{mz_2 + nz_1}{m + n} \right)$$

Similarly if  $R$  divides  $PQ$  externally in the ratio of  $m : n$ , then the coordinates of  $R$  are

$$\left( \frac{mx_2 - nx_1}{m - n}, \frac{my_2 - ny_1}{m - n}, \frac{mz_2 - nz_1}{m - n} \right)$$

### Illustration 3

**Question:** Find the ratio in which  $yz$ -plane divides the line joining  $(2, 4, 5)$  and  $(3, 5, 7)$ .

**Solution:** Let the ratio be  $\lambda : 1$

$$\text{x-coordinate is } \frac{3\lambda + 2}{\lambda + 1} = 0 \Rightarrow \lambda = -\frac{2}{3}$$

\(\therefore\) 2 : 3 (externally)

### Illustration 4

**Question:** Show that the points  $(2, -3, 4)$ ,  $(-1, 2, 1)$  and  $(0, \frac{1}{3}, 2)$  are collinear and find the ratio in which the third point divides the line joining first two.

**Solution:** Let  $A$  denote the point  $(2, -3, 4)$  and  $B$  the point  $(-1, 2, 1)$

Let  $C$  divides the line joining  $A$  and  $B$  in the ratio of  $\lambda : 1$

\(\therefore\) co-ordinates of  $C$  are

$$\left( \frac{2 - \lambda}{\lambda + 1}, \frac{-3 + 2\lambda}{\lambda + 1}, \frac{4 + \lambda}{\lambda + 1} \right) \equiv \left( 0, \frac{1}{3}, 2 \right) \text{ \{given\}}$$

$$\therefore \frac{2 - \lambda}{\lambda + 1} = 0 \Rightarrow \lambda = 2$$

$$\text{and } \frac{-3 + 2\lambda}{\lambda + 1} = \frac{1}{3}$$

$$\Rightarrow -9 + 6\lambda = \lambda + 1$$

$$\Rightarrow 5\lambda = 10 \quad \Rightarrow \quad \lambda = 2$$

$$\text{Also } \frac{4 + \lambda}{\lambda + 1} = 2$$

$$\Rightarrow 4 + \lambda = 2\lambda + 2$$

$$\Rightarrow \lambda = 2$$

From each equation we get the value of  $\lambda$  as 2.

$\therefore$  The point  $A$ ,  $C$  and  $B$  are collinear.

$\therefore$  The point  $C$  divides the line joining  $A$  and  $B$  internally in the ratio of 2 : 1.

---

### PRACTICE PROBLEMS

- PP1.** Show that the points  $(a, b, c)$ ,  $(b, c, a)$  and  $(c, a, b)$  are the vertices of an equilateral triangle.
- PP2.** Find the ratio in which the point  $C(5, 9, -14)$  divides the join of  $A(2, -3, 4)$ ,  $B(3, 1, -2)$ .
- PP3.** Find the coordinates of the point which divides the line segment joining the points  $(1, -2, 3)$  and  $(3, 4, -5)$  in the ratio 2 : 3 (i) internally and (ii) externally.
- PP4.** Find the coordinates of the centroid of the triangle whose vertices are  $(x_1, y_1, z_1)$ ,  $(x_2, y_2, z_2)$  and  $(x_3, y_3, z_3)$ .
- PP5.** Find the distance of the point  $P(a, b, c)$  from  $x$ -axis.
- PP6.** Find the area of the triangle formed by joining the points whose coordinates are  $(1, 0, 0)$ ,  $(1, 2, 0)$  and  $(1, 0, 3)$  in sq. units.
-


SOLVED SUBJECTIVE EXAMPLES

**Example 1:**

Find the co-ordinate of the point  $P$  which is five sixth of the way from  $A(-2, 0, 6)$  to  $B(10, -6, -12)$ .

**Solution :**

Let  $P(x, y, z)$  be the required point



$\Rightarrow P$  divides  $AB$  in ratio  $5 : 1$

$$\left[ \because AP = \frac{5}{6} AB \Rightarrow \frac{AP}{AB} = \frac{5}{6} \Rightarrow \frac{AP}{AB - AP} = \frac{5}{6 - 5} \Rightarrow \frac{AP}{PB} = \frac{5}{1} \right]$$

$$\Rightarrow (x, y, z) = \left( \frac{5 \times 10 + 1 \times -2}{5 + 1}, \frac{5 \times -6 + 1 \times 0}{5 + 1}, \frac{5 \times -12 + 1 \times 6}{5 + 1} \right)$$

$\Rightarrow (x, y, z) = (8, -5, -9) \Rightarrow P(8, -5, -9)$  is the required point.

**Example 2:**

Prove that the line through  $A(0, 1, 1)$ ,  $B(4, 5, 1)$  intersects the line through  $C(3, 9, 4)$  and  $D(4, 4, 4)$ .


**Solution :**

The lines  $AB$  and  $CD$  intersect if there exists a common point  $P$  dividing  $AB$  in some ratio say  $\lambda : 1$  and  $CD$  in some ratio say  $\mu : 1$  where  $\lambda + 1 \neq 0, \mu + 1 \neq 0$

$$\text{i.e., if } \left( \frac{4\lambda}{\lambda + 1}, \frac{5\lambda - 1}{\lambda + 1}, \frac{\lambda - 1}{\lambda + 1} \right) = \left( \frac{-4\mu + 3}{\mu + 1}, \frac{4\mu + 9}{\mu + 1}, \frac{4\mu + 4}{\mu + 1} \right)$$

$$\text{if } \begin{cases} \frac{4\lambda}{\lambda + 1} = \frac{-4\mu + 3}{\mu + 1} \\ \frac{5\lambda + 1}{\lambda + 1} = \frac{4\mu + 9}{\mu + 1} \\ \frac{\lambda - 1}{\lambda + 1} = \frac{4\mu + 4}{\mu + 1} \end{cases} \text{ are consistent.}$$

$$\text{i.e., } \lambda = \frac{-5}{3}$$



$$\text{i.e., if } \frac{4\left(-\frac{5}{3}\right)}{\frac{-5}{3}+1} = \frac{-4\mu+3}{\mu+1}, \frac{5\left(-\frac{5}{3}\right)-1}{\frac{-5}{3}+1} = \frac{4\mu+9}{\mu+1}, \lambda = -\frac{5}{3} \text{ are consistent.}$$

$$\text{i.e., if } \frac{-20}{-2} = \frac{-4\mu+3}{\mu+1}, \frac{-28}{-2} = \frac{4\mu+9}{\mu+1}, \lambda = -\frac{5}{3} \text{ are consistent.}$$

$$\text{i.e., if } 10 = \frac{-4\mu+3}{\mu+1}, 14 = \frac{4\mu+9}{\mu+1}, \lambda = -\frac{5}{3} \text{ are consistent.}$$

$$\text{i.e., if } 14\mu = -7, 10\mu = -5, \lambda = -\frac{5}{3} \text{ are consistent.}$$

$$\text{i.e., if } \mu = -\frac{1}{2}, \mu = -\frac{1}{2}, \lambda = -\frac{5}{3} \text{ which is true.}$$

$\therefore$  Lines  $AB$  and  $CD$  intersect in  $P$ .

### Example 3:

Using section formula, prove that the three points  $Q(2, 3, 5)$ ,  $R(1, 2, 3)$  and  $S(7, 0, -1)$  are collinear. Also find the ratio in which the third point divides the segment joining first two points.

### Solution :

Let  $A(-2, 3, 5)$ ,  $B(1, 2, 3)$ ,  $C(7, 0, -1)$  be the given points.

The point which divides  $AB$  in the ratio  $k : 1$  is  $\left(\frac{k-2}{k+1}, \frac{2k+3}{k+1}, \frac{3k+5}{k+1}\right)$  ... (i)

If this is the point  $C(7, 0, -1)$ , then  $\frac{k-2}{k+1} = 7, \frac{2k+3}{k+1} = 0, \frac{3k+5}{k+1} = -1$

First equation gives  $k-2 = 7k+7$

$$\text{i.e., } 6k = -9, \therefore k = -\frac{9}{6} = -\frac{3}{2}$$

Second equation gives  $2k+3 = 0, \therefore k = -\frac{3}{2}$

Third equation gives  $3k+5 = -k-1$  or  $4k = -6$

$$\therefore k = -\frac{6}{4} = -\frac{3}{2}$$

Hence for  $k = -\frac{3}{2}$  the point given (i) coincides with  $C$ .

$\therefore C$  divides  $AB$  externally in the ratio  $3 : 2$

Thus  $A, B, C$  are collinear.

**Example 4:**

Find the ratio in which the surface  $x^2 + y^2 + z^2 = 25$  divides the line segment joining  $(0, 1, 2)$  and  $(3, 4, 5)$ .

**Solution :**

Given surface is  $x^2 + y^2 + z^2 = 25$

Let  $A \equiv (0, 1, 2)$ ,  $B \equiv (3, 4, 5)$  ... (i)

Let surface (i) divide the line segment  $AB$  internally in the ratio  $k : 1$  at  $P$ .

$$\text{Then } P = \left( \frac{3k}{k+1}, \frac{4k+1}{k+1}, \frac{5k+2}{k+1} \right)$$

Since  $P$  lies on surface (i),

$$\therefore \frac{9k^2 + (4k+1)^2 + (5k+2)^2}{(k+1)^2} = 25 \text{ or } 50k^2 + 28k + 5 = 25(k^2 + 2k + 1)$$

$$\text{or } 25k^2 - 23k - 20 = 0 \text{ or } k = \frac{23 \pm \sqrt{2599}}{50} = \frac{23 + \sqrt{2529}}{50}, \frac{23 - \sqrt{2529}}{50}$$

Hence surface (i) divides the line segment  $AB$  internally in the ratio  $\frac{23 + \sqrt{2529}}{50}$  and externally in the ratio  $\frac{23 - \sqrt{2599}}{50}$ .


**Example 5:**

Find the locus of a locus of a point while moves so that its distances from the points  $(3, 4, -5)$  and  $(-2, 1, 4)$ .

**Solution :**

Let  $A = (3, 4, -5)$ ,  $B = (-2, 1, 4)$

Let  $P(x, y, z)$  be any point on the locus.



$\therefore$  By the given conditions.  $PA = PB$

$$\therefore \sqrt{(x-3)^2 + (y-4)^2 + (z+5)^2} = \sqrt{(x+2)^2 + (y-1)^2 + (z-4)^2}$$

$$\Rightarrow x^2 + 9 - 6x + y^2 + 16 - 8y + z^2 + 25 + 10z$$

$$= x^2 + 4 + 4x + y^2 + 1 - 2y + z^2 + 16 - 8z$$

$$\Rightarrow -10x - 6y + 18z = 29$$



$$\Rightarrow 10x + 6y - 18z + 29 = 0$$

**Example 6:**

Find the equation of the set of points  $P$ , the sum of whose distance from  $A(4, 0, 0)$  and  $B(-4, 0, 0)$  is equal to 10.

**Solution :**

Given points are  $A(4, 0, 0)$  and  $B(-4, 0, 0)$

Let  $P(x, y, z)$  be any point such that  $|PA| + |PB| = 10$

$$\Rightarrow \sqrt{(x-4)^2 + (y-0)^2 + (z-0)^2} + \sqrt{(x+4)^2 + (y-0)^2 + (z-0)^2} = 10$$

$$\Rightarrow \sqrt{(x+4)^2 + y^2 + z^2} = 10 - \sqrt{(x-4)^2 + y^2 + z^2}$$

Squaring the two sides, we get

$$(x+4)^2 + y^2 + z^2 = 100 + (x-4)^2 + y^2 + z^2 - 20\sqrt{(x-4)^2 + y^2 + z^2}$$

$$\text{or } (x+4)^2 - (x-4)^2 - 100 = -20\sqrt{(x-4)^2 + y^2 + z^2}$$

$$\text{or } 16x - 100 = -20\sqrt{(x-4)^2 + y^2 + z^2}$$

Dividing by  $-4$ , we obtain

$$-4x + 25 = 5\sqrt{(x-4)^2 + y^2 + z^2}$$

$$\text{Squaring, we get } 16x^2 + 625 - 200x = 25\{x^2 - 8x + 16 + y^2 + z^2\}$$

$$\text{or } 9x^2 + 25y^2 + 25z^2 = 225 \quad \text{or } \frac{9x^2}{225} + \frac{25y^2}{225} + \frac{25z^2}{225} = 1$$

$$\text{or } \frac{x^2}{25} + \frac{y^2}{9} + \frac{z^2}{9} = 1, \text{ which is the required equation.}$$

**EXERCISE – I**

1. Name the octants in which the following points lie:  
(1, 2, 3), (4, -2, -5), (-3, 2, -5) and (-4, -6, 7).
2. Find the distance between the points  $A(5, 1, 2)$  and  $B(4, 6, -1)$ .
3. Show that  $A(-2, 3, 5)$ ,  $B(1, 2, 3)$  and  $C(7, 0, -1)$  are collinear.
4. Determine the values of  $a$  and  $b$  so that the points  $(a, b, 3)$ ,  $(2, 0, -1)$  and  $(1, -1, -3)$  are collinear.
5. Let  $A(2, 1, -3)$  and  $B(5, -8, 3)$  be two given points. Find the coordinates of the points of trisection of the line segment  $AB$ .
6. Show that the points  $A(1, 1, 1)$ ,  $B(-2, 4, 1)$ ,  $C(-1, 5, 5)$  and  $D(2, 2, 5)$  are the vertices of a square.
7. Find the coordinates of the point where the line joining  $A(3, 4, 1)$  and  $B(5, 1, 6)$  crosses the  $xy$ -plane.
8. If the three consecutive vertices of a parallelogram be  $A(3, 4, -1)$ ,  $B(7, 10, -3)$  and  $C(5, -2, 7)$ , find the fourth vertex  $D$ .
9. The midpoints of the sides of a triangle are  $(1, 5, -1)$ ,  $(0, 4, -2)$  and  $(2, 3, 4)$ . Find its vertices.
10. Find the point in  $yz$ -plane which is equidistant from the points  $A(3, 2, -1)$ ,  $B(1, -1, 0)$  and  $C(2, 1, 2)$ .

**EXERCISE – II**

1. Planes are drawn parallel to the co-ordinate planes through the points  $(1, 2, 3)$  and  $(3, -4, -5)$ . Find the length of diagonal of parallelepiped so formed.
2. Show that the line segment joining the points  $A(1, 2, 3)$ ,  $B(4, 5, 6)$  is parallel to the line segment joining the points  $C(-4, 3, -6)$  and  $D(2, 9, 0)$ .
3. Find the length of medians of the triangle having vertices  $A(0, 0, 6)$ ,  $B(0, 4, 0)$  and  $C(6, 0, 0)$ .
4. Find the coordinates of a point on  $y$ -axis which are at a distance of  $5\sqrt{2}$  from the point  $P(3, -2, 5)$ .
5. The centroid of a triangle  $ABC$  is at the point  $(1, 1, 1)$ . If the coordinates of  $A$  and  $B$  are  $(3, -5, 7)$  and  $(-1, 7, -6)$  respectively, find the coordinates of the point  $C$ .
6. Let  $A, B, C$  be the feet of perpendiculars drawn from a point  $P$  to  $x, y$  and  $z$  axes respectively. Find the co-ordinates of  $A, B, C$ ; if co-ordinates of  $P$  are  
(i)  $(3, 1, 2)$                       (ii)  $(3, -6, 2)$
7. Find the equation of the set of points  $P$ , the sum of whose distances from  $A(4, 0, 0)$  and  $B(-4, 0, 0)$  is equal to 10.
8. Find the locus of a point  $P$  which moves in such a way that  $2PA = 3PB$ , where  $A(-2, 1, 3)$ , and  $B(3, -1, 2)$  are the given points.
9. Find the equation of set of points  $P$  such that  $PA^2 + PB^2 = 2k^2$ , where  $A$  and  $B$  are the points  $(3, 4, 5)$  and  $(-1, 3, -7)$  respectively.
10. The centroid of a triangle  $ABC$  is at the point  $(1, 1, 1)$ . If the coordinates of  $A$  and  $B$  are  $(3, -5, 7)$  and  $(-1, 7, -6)$  respectively, find the coordinates of the point  $C$ .

**ANSWERS****ANSWERS TO PRACTICE PROBLEMS**

PP2. (3 : 2 externally)

PP3. (-3, -14, 19)

PP4.  $\left(\frac{\sum x_1}{3}, \frac{\sum y_1}{3}, \frac{\sum z_1}{3}\right)$

PP5.  $\sqrt{b^2 + c^2}$

PP6. 3

**ANSWERS TO EXERCISE – I**

1. I, VIII, VI, III
2.  $\sqrt{35}$  units
4.  $a = 4, b = 2$
5.  $(3, -2, -1), (4, -5, 1)$
7.  $\left(\frac{13}{5}, \frac{23}{5}, 0\right)$
8.  $(1, -8, 9)$
9.  $(1, 2, 3) (3, 4, 5) (-1, 6, -7)$
10.  $\left(0, \frac{31}{16}, \frac{-3}{16}\right)$

**ANSWERS TO EXERCISE – II**

1.  $\sqrt{104}$
3.  $7, \sqrt{34}, 7$
4.  $(0, 2, 0)$  and  $(0, -6, 0)$
5.  $(1, 1, 2)$
6. (i)  $A(3, 0, 0), B(0, 1, 0), C(0, 0, 2)$   
(ii)  $A(3, 0, 0), B(0, -6, 0), C(0, 0, 2)$
7.  $9x^2 + 25y^2 + 25z^2 = 225$
8.  $5(x^2 + y^2 + z^2) - 70x + 26y - 12z + 70 = 0$
9.  $2x^2 + 2y^2 + 2z^2 - 4x - 14y + 4z = 2k^2 - 109$
10.  $(1, 1, 2)$